



### Please take a minute...

## to learn about my diagnosis of Parkinson's disease

and critical information regarding my inpatient care.



- I need my PD medications on time, every time!
- My PD symptoms may fluctuate throughout the day.
- Dopamine blocking drugs, such as haloperidol (Haldol) and metoclopramide (Reglan), can worsen my condition and should be avoided.
- Quetiapine (Seroquel) and clozapine (Clozaril) are the only antipsychotic medications considered safe for PD.
- Range of motion exercises and physical therapy are important to avoid excess rigidity.
- Fall precautions should be followed.
- Contact my PD provider with questions/concerns and prior to adjusting my PD treatment regimen at: \_\_\_\_\_

For more information on PD, go to [www.parkinsons.va.gov](http://www.parkinsons.va.gov)



### Please take a minute...

## to learn about my diagnosis of Parkinson's disease

and critical information regarding my inpatient care.



- I need my PD medications on time, every time!
- My PD symptoms may fluctuate throughout the day.
- Dopamine blocking drugs, such as haloperidol (Haldol) and metoclopramide (Reglan), can worsen my condition and should be avoided.
- Quetiapine (Seroquel) and clozapine (Clozaril) are the only antipsychotic medications considered safe for PD.
- Range of motion exercises and physical therapy are important to avoid excess rigidity.
- Fall precautions should be followed.
- Contact my PD provider with questions/concerns and prior to adjusting my PD treatment regimen at : \_\_\_\_\_

For more information on PD, go to [www.parkinsons.va.gov](http://www.parkinsons.va.gov)



### Please take a minute...

## to learn about my diagnosis of Parkinson's disease

and critical information regarding my inpatient care.



- I need my PD medications on time, every time!
- My PD symptoms may fluctuate throughout the day.
- Dopamine blocking drugs, such as haloperidol (Haldol) and metoclopramide (Reglan), can worsen my condition and should be avoided.
- Quetiapine (Seroquel) and clozapine (Clozaril) are the only antipsychotic medications considered safe for PD.
- Range of motion exercises and physical therapy are important to avoid excess rigidity.
- Fall precautions should be followed.
- Contact my PD provider with questions/concerns and prior to adjusting my PD treatment regimen at: \_\_\_\_\_

For more information on PD, go to [www.parkinsons.va.gov](http://www.parkinsons.va.gov)



### Please take a minute...

## to learn about my diagnosis of Parkinson's disease

and critical information regarding my inpatient care.



- I need my PD medications on time, every time!
- My PD symptoms may fluctuate throughout the day.
- Dopamine blocking drugs, such as haloperidol (Haldol) and metoclopramide (Reglan), can worsen my condition and should be avoided.
- Quetiapine (Seroquel) and clozapine (Clozaril) are the only antipsychotic medications considered safe for PD.
- Range of motion exercises and physical therapy are important to avoid excess rigidity.
- Fall precautions should be followed.
- Contact my PD provider with questions/concerns and prior to adjusting my PD treatment regimen at: \_\_\_\_\_

For more information on PD, go to [www.parkinsons.va.gov](http://www.parkinsons.va.gov)